

Soy intake is associated with lowering blood pressure in adults: A systematic review and meta-analysis of randomized double-blind placebo-controlled trials

Mosallanezhad, Z., Ranjbar, S., Mahmoodi, M., Hosseini, R., Clark, C. C. T., Carson-Chahhoud, K., Norouzi, Z., Abbasian, A. & Jalali, M.

Published PDF deposited in Coventry University's Repository

Original citation:

Mosallanezhad, Z, Ranjbar, S, Mahmoodi, M, Hosseini, R, Clark, CCT, Carson-Chahhoud, K, Norouzi, Z, Abbasian, A & Jalali, M 2021, 'Soy intake is associated with lowering blood pressure in adults: A systematic review and meta-analysis of randomized double-blind placebo-controlled trials', *Complementary Therapies in Medicine*, vol. 59, 102692.

<https://dx.doi.org/10.1016/j.ctim.2021.102692>

DOI 10.1016/j.ctim.2021.102692

ISSN 0965-2299

Publisher: Elsevier

This is an open access article under the CC BY-NC-ND license



Soy intake is associated with lowering blood pressure in adults: A systematic review and meta-analysis of randomized double-blind placebo-controlled trials

Zahra Mosallanezhad^{a,b,1}, Marzieh Mahmoodi^{a,b,1}, Sara Ranjbar^{a,b}, Razieh Hosseini^c, Cain C. T. Clark^d, Kristin Carson-Chahhoud^{e,f}, Zahra Norouzi^g, Ali Abbasian^h, Zahra Sohrabiⁱ, Mohammad Jalali^{a,*}

^a Nutrition Research Center, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

^b Student Research Committee, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

^c Student Research Committee, Department of Nutrition, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

^d Centre for Sport, Exercise and Life Sciences, Coventry University, Coventry CV15FB, UK

^e Australian Centre for Precision Health, School of Health Sciences, University of South Australia, Australia

^f School of Medicine, the University of Adelaide, South Australia, Australia

^g Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran

^h Department of Physical Education & Sport Sciences, Masjed-Soleiman Branch, Islamic Azad University, Masjed-Soleiman, Iran

ⁱ Department of Community Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

ARTICLE INFO

Keywords:

Soy
Blood pressure
Hypertension
Review
Meta-analysis

ABSTRACT

Background: Soy has several beneficial effects on cardiovascular disease (CVD); however, results of clinical trial studies are equivocal. Thus, the present study sought to discern the efficacy of soy intake on blood pressure.

Methods: The search process was conducted in PubMed, Scopus, Web of Science, and Cochrane Library, to ascertain studies investigating the efficacy of soy intake on blood pressure in adults, published up to June 2020. A random-effects model was applied to pool mean difference and 95 % confidence interval (CI). Begg's and Egger's methods were conducted to assess publication bias.

Results: Pooled effects from 17 effect sizes revealed a significant improvement in systolic blood pressure (SBP) (-1.70 ; -3.34 to -0.06 mmHg; $I^2 = 45.4$ %) and diastolic blood pressure (DBP) (-1.27 ; -2.36 to -0.19 mmHg, $I^2 = 43.9$ %) following soy consumption, in comparison with controls. Subgroup analysis demonstrated a reduction in both SBP and DBP in younger participants with lower baseline DBP and intervention durations of <16 weeks.

Conclusion: Our results suggest that soy intake is associated with an ameliorating effect on blood pressure in adults.

1. Introduction

Soy is a traditional food that is globally popular, especially in Asia, and is widely used to produce various food products, such as soybean oil, soy milk, soy flour, and many retail food products.¹ Utilizing soy products in the food processing industry makes it possible to achieve significant economic effects by reducing production costs and standardizing quality.^{2,3} Soybean contains compounds such as protein, fiber, vitamins,

minerals, and phytochemicals, which has led to considerable research interest regarding its effects on various diseases.^{1,4} Accordingly, previous studies have reported protective features of soy in diabetes, cancer, osteoporosis, and menopausal problems,³ with further studies confirming its beneficial effects on cardiovascular disease (CVD).⁵

One of the most important risk factors of CVD is hypertension, which represents one of the most prevalent non-communicable diseases, worldwide.⁶ Several strategies, including medication and lifestyle

* Corresponding author at: Nutrition research Center, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Razi Ave, Shiraz, 7153675541, Iran.

E-mail address: Mj.artak@gmail.com (M. Jalali).

¹ These authors contributed equally to the work and should be regarded as equal first authors.

<https://doi.org/10.1016/j.ctim.2021.102692>

Received 17 January 2021; Accepted 20 February 2021

Available online 24 February 2021

0965-2299/© 2021 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. The process of study selection.

modification are recommended to manage the disorder,⁷ whilst numerous studies have been performed to investigate the effects of nutrients on blood pressure.⁸ For instance, the Dietary Approaches to Stop Hypertension (DASH) diet has been advocated as part of the treatment protocol for hypertensive patients.⁹ Moreover, there are several studies demonstrating the effectiveness of macronutrients on blood pressure.^{10,11} In the case of total protein intake, however, the results manifest in the literature are contradictory, although the positive effect of plant based protein on blood pressure has been confirmed.¹² In addition, although observational studies^{13,14} have confirmed the efficacy of soy as a vegetable protein on blood pressure, the results of clinical trial studies are equivocal.^{15,16}

Due to the limited number of participants in previous clinical trial studies,^{17,18} as well as differences in intervention duration,^{19,20} age,^{21,22} and baseline blood pressure,^{18,21,23} it is not possible to draw any firm conclusion regarding the effect of soy on blood pressure. Therefore, in this study we aimed to identify the effect of soy intake on blood pressure by performing a systematic review and meta-analysis of randomized clinical trials.

2. Methods

We undertook our study in accordance with guidelines proposed by the *Preferred Reporting Items for Systematic reviews and Meta-Analyses* (PRISMA).²⁴ The review protocol was registered with PROSPERO (CRD42020214728).

2.1. Search process

To identify relevant papers investigating the effect of soy on blood pressure, we undertook a systematic search in PubMed, Embase, Scopus, Web of Science, and Cochrane Library, for all available publications up to June 2020, using a combination the relevant terms in titles and abstracts, without making any restrictions (Supplementary material 1). In some cases, we used the wildcard term “*” to increase sensitivity and checked the reference lists of the relevant reviews and Google Scholar to supplement the database search strategy, which was prepared with in-group consensus. After loading retrieved records into EndNote, duplicates were removed.

2.2. Inclusion and exclusion criteria

Four authors independently assessed the records retrieved from the database search. Randomized controlled trials (RCTs) investigating the association of soy intake and blood pressure in adults (≥ 18 years of age) were identified for inclusion. To maximize methodological rigor, only double-blind placebo-controlled trials were then shortlisted for inclusion, to increase quality of evidence, and reduce potential bias in the meta-analysis. Review articles, dissertations, brief reports, observation-designed and non-English language studies were excluded, as were those lacking clinical information or statistical data required for analysis (e.g. baseline of blood pressure, standard deviation (SD), standard error (SE), 95 % confidence intervals (CIs) and interquartile (IQR)). Disagreements in the inclusion or exclusion of any study were resolved by group discussion. For example, one study did not clearly mention term “placebo”

Table 1
Characteristics of the included RCTs.

Study	Country	Type of study	Population	Age group (C vs. I)	Body weight (C vs. I)	BMI (C vs. I)	Intervention type	Sample size	Duration (weeks)	Baseline SBP (mmHg)	Baseline DBP (mmHg)
Teede (2001)	Australia	Randomized, double-blind, placebo-controlled trial	Men and Postmenopausal Women	60 vs. 61	74 vs. 72	26 vs. 25	Soy protein	179	12	130	76
Sagara (2004)	Scotland	Randomized, double-blind, placebo-controlled trial	Hypercholesterolemic and / or hypertensive	52.2 vs. 52.2	83.8 vs. 85.1	27.2 vs. 27.6	Soy protein	50	5	142	87.1
He (2005)	China	Randomized, double-blind, controlled trial.	Healthy	51.4 vs. 50.8	70.6 vs. 70	26.8 vs. 26.9	Soybean protein	276	12	134.7	84.7
Kim (2005)	Korea	Randomized, double-blind, placebo-controlled trial	T2DM	61.7 vs. 59.9	61.1 vs. 64.1	23.8 vs. 24.4	Soybean	30	13	137.5	86.6
Hermansen (2005)	Denmark	Randomized, double-blind, placebo-controlled trial	Hypercholesterolemic	58 vs. 60.6	75.3 vs. 77.1	25.6 vs. 26.4	Soy protein	100	24	133	80.2
Aubertin-Leheudre (2008)	Canada	Randomized, double-blind, placebo-controlled trial	Obese Postmenopausal Women	57.7 vs. 57.1	82.5 vs. 79.6	32.8 vs. 31.2	Soy isoflavone	39	25	125.1	79
Chan (2008)	China	Randomized, double-blind, placebo-controlled trial	Ischaemic stroke	65.8 vs. 66.8	NR*	25 vs. 26.2	Soy isoflavone	102	12	141	77
Kwak (2010)	Korea	Randomized, double-blind, placebo-controlled trial	Prediabetes and newly diagnosed T2DM	57.6 vs. 56.8	65.8 vs. 62.6	24.8 vs. 24.1	Black soy peptide	42	12	125.1	73.6
Wong (2012)	Netherlands	Randomized, double-blind, placebo-controlled trial	Menopausal women with high or normal blood pressure	55.5 vs. 55.8	68.9 vs. 67.6	25.4 vs. 25.3	Soy hypocotyl isoflavones	24	6	140.1	82.8
Liu 1 (2013)	China	Randomized, double-blind, placebo-controlled trial	Postmenopausal women with mild hyperglycemia	54.8 vs. 54.3	54.4 vs. 56.1	22.6 vs. 23.6	Soy protein and isoflavones	180	25	127.9	77
Liu 2 (2013)	China	Randomized, double-blind, placebo-controlled trial	Postmenopausal women with mild hyperglycemia	54.8 vs. 56.3	54.4 vs. 56.8	22.6 vs. 23.9	Soy isoflavones	180	25	125.9	78.2
Kim (2013)	Korea	Randomized, double-blind, placebo-controlled trial	Postmenopausal women	53.5 vs. 53.7	56.5 vs. 57.7	23.3 vs. 23.2	Soy isoflavone	85	12	116.1	74.6
Squadrito (2013)	Italy	Randomized, double-blind, placebo-controlled trial	Postmenopausal women with metabolic syndrome	55.4 vs. 55.6	NR	31.8 vs. 31.8	Soy isoflavone	108	51	135.7	78.7
Cheng (2013)	Taiwan	Randomized, double-blind, placebo-controlled trial	Healthy postmenopausal women	56.1 vs. 57	55.1 vs. 55.1	22.9 vs. 23.1	Soy isoflavone	82	52	119	77.3
Husain (2015)	Iran	Randomized, double-blind, placebo-controlled trial	Postmenopausal women	50.32 vs. 50.93	NR	NR	Soy	61	8	125.77	81.77
Liu 1 (2015)	China	Randomized, double-blind, placebo-controlled trial	Postmenopausal women with prehypertension	58.5 vs. 57.6	57.6 vs. 56.5	NR	Soy	180	24	130.7	81.6
Liu 2 (2015)	China	Randomized, double-blind, placebo-controlled trial	Postmenopausal women with prehypertension	58.5 vs. 57.7	57.6 vs. 56.5	NR	Soy isoflavone	180	24	131.9	81.6

C (control group); I (intervention group); BMI (body mass index); T2DM (type 2 diabetes mellitus), NR (not reported).

Table 2

The methodological quality of included RCTs on effect of soy intake on blood pressure based on review authors' judgments about each risk of bias item for each included study.

Study	Sequence generation	Allocation concealment	Blinding	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other bias
Teede (2001)							
Sagara (2004)							
He (2005)							
Kim (2005)							
Hermansen (2005)							
Aubertin-Leheudre (2008)							
Chan (2008)							
Kwak (2010)							
Wong (2012)							
Liu (2013)							
Kim (2013)							
Squadrito (2013)							
Cheng (2013)							
Husain (2015)							
Liu (2015)							

as the control,²¹ however, the study-design was comparable to a placebo-controlled design. Following discussion with clinical and statistical experts, this study was included. In addition, studies were excluded if the control group was not clinically comparable with the intervention, and therefore inclusion would have biased the results. Trials that prescribed effective nutrients, diet, medicines, or other combinations concomitant to soy, in comparison with controls were also excluded.

2.3. Data extraction

After identifying studies for inclusion, a combination of four authors extracted clinical and statistical data, including: first authors last name, corresponding author's e-mail, publication year, country, population, participants' characteristics, intervention and control type, dose and type of supplement, treatment duration, sample size, methodological quality, mean change of interested outcome from baseline at the end of trial, related SD, SE, 95 % CI or IQR. Where required appropriate transformations were undertaken to calculate the SD from studies reporting: SE ($SD = SE \times \sqrt{n}$), 95 % CI ($SD = \sqrt{n \times (upper\ limit - lower\ limit) / 3.92}$) or IQR ($SD = IQR / 1.35$).²⁵ In addition, WebPlotDigitizer (<https://automeris.io/WebPlotDigitizer>) was used to derive the mean change and the corresponding SD when the data were plotted, but not reported. In the case of an article with two different intervention groups

and one control group, the intervention groups were considered as different studies, and to avoid giving more weight to them; number of participants in the control group was divided by 2.

2.4. Quality appraisal

Two authors independently assessed quality in the included studies using the Cochrane Collaboration risk of bias assessment tool,²⁶ which is scored across seven domains being: random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective reporting (reporting bias), and other sources of bias. Each included study also received an overall grading of Good (> 2 low risk domains), Fair (= 2 low risk domains), or Poor (< 2 low risk domains).

2.5. Statistical analysis

Weighted mean difference (WMD) and 95 % CI was calculated in Stata v.13, applying a random-effects method. Inter-study heterogeneity was evaluated by checking I^2 index (low: <50%, high: >50%).²⁷ Sub-group analyses were planned by effective clinical covariates i.e. age (> 56, < 56 years old), baseline blood pressure (SBP: > 130, < 130 mmHg; DBP: > 80, < 80 mmHg), and intervention duration (< 16, > 16 weeks).




Fig. 2. Forest plot detailing WMDs and 95 % CIs for the meta-analyses of SBP (A) and DBP (B).




Fig. 3. Findings from subgroup analysis based on age (< 56, > 56 years old) (A), baseline blood pressure (< 130, > 130 mmHg) (B), duration (< 16, > 16 weeks) (C) and region (Asia, Europe, America) (D) regarding the effect of soy consumption on SBP.

Mean changes and their SDs of relevant outcomes were obtained by using the following equations: $[\text{mean-post} - \text{mean-baseline}]$ and $[(\sqrt{((\text{SD pre})^2 + (\text{SD post})^2) - [2r \times \text{SD pre} \times \text{SD post}]})]$,^{28,49,50} respectively. In addition, a sensitivity analysis was applied by discarding each trial in turn, to ensure robustness of results. A Begg's rank-correlation²⁹ and Egger's regression asymmetry³⁰ were performed to evaluate potential publication bias. *P*-values of less than 0.05 represented statistical significance.

3. Results

3.1. Results of the search and study characteristics

From 3163 records, 15 RCTs with 17 effect sizes were identified for inclusion, all of which were able to be included in the meta-analysis (Fig. 1). These studies were published between 2001 and 2015. Ten were conducted in Asia,^{17,19,20,21,22,31,32,33,34,35} four in Europe,^{18,23,36,37} and another one in the United States of America.³⁸ Participant age ranged from 50.32–66.8 years. Intervention durations were less than 16 weeks^{17,18,20,21,22,23,31,32,33} or more than 16 weeks^{19,34,35,36,37,38} in nine and six trials, respectively (Table 1).

Baseline blood pressure across studies was reported as SBP < 130

mmHg^{17,20,33,34,35,38} and > 130 mmHg^{18,19,21,22,23,31,32,36,37} in six and nine RCTs, respectively. In addition, enrolled participants of seven and eight studies had > 80 mmHg^{18,19,20,21,23,32,36} and < 80 mmHg^{17,22,31,33,34,35,37} DBP, respectively.

3.2. Quality assessment for included studies

Results of the quality assessment are shown in Table 2. Overall, study quality was assessed as good, with a low risk of bias the dominant classification across all domains. All studies were assessed as having low risk of bias for sequence generation and blinding of participants. Six studies were assessed as having unclear risk of bias for allocation concealment, five had unclear risk for blinding of outcome assessment, one had unclear risk for selective reporting and three had unclear risk for other potential sources of bias. Two studies were assessed as having high risk of bias for incomplete outcome data, with three assessed as unclear and the remaining 10 studies as low risk.

3.3. Meta-analysis of blood pressure outcomes

A significant reduction in both SBP (WMD = -1.70 mmHg, 95 % CI = [-3.34, -0.06], *P* = 0.04, *I*² = 45.4 %) and DBP (WMD = -1.27




Fig. 4. Findings from subgroup analysis based on age (< 56, > 56 years old) (A), baseline blood pressure (< 130, > 130 mmHg) (B), duration (< 16, > 16 weeks) (C) and region (Asia, Europe, America) (D) regarding the effect of soy consumption on DBP.

mmHg, 95 % CI = [-2.36, -0.19], $P = 0.02$, $I^2 = 43.9\%$) were observed in the soy intake group when compared with controls (Fig. 2).

Subgroup analyses, identified a greater reduction in blood pressure for participants aged younger than 56 years (Fig. 3, A and Fig. 4, A) and lower baseline diastolic blood pressure (Fig. 4, B). In addition, both systolic (Fig. 3, C) and diastolic (Fig. 4, C) blood pressure were significantly reduced when soy intake duration was < 16 weeks.

3.4. Publication bias and sensitivity analysis

Egger's regression and Begg's rank-correlation tests indicated that there was no significant publication bias for SBP (P for Egger's test = 0.71 and Begg's test = 0.99) (Fig. 5, A) or DBP (P for Egger's test = 0.94 and Begg's test = 0.51) (Fig. 5, C). According to the findings from sensitivity analysis, pooled effect sizes obtained for the effect of soy intake on SBP (Fig. 5, B) and DBP (Fig. 5, D) were not sensitive to any particular study or group of studies.

4. Discussion

Data from 17 effect sizes were available to examine efficacy of soy consumption on blood pressure in adults. Our results showed that soy

consumption significantly improves SBP and DBP. Subgroup analyses identified a greater reduction in blood pressure among younger participants with lower baseline diastolic blood pressure, and in trials lasting for < 16 weeks duration.

The effect of soy consumption on blood pressure has been previously assessed in several RCTs. A study by Washburn et al. indicated that 40 g of soy protein containing 68 mg of phytoestrogens, improved blood pressure abnormality.³⁹ In Teede et al., it was demonstrated that administration of 40 g of soy protein containing 118 mg of isoflavones, improved blood pressure in healthy men and women.³¹ Furthermore, Welty et al. showed that a soy nut diet (containing 25 g of soy protein and 101 mg of aglycone isoflavones) lowered systolic blood pressure in hypertensive and normotensive patients⁴⁰; whilst, Rivas et al. reported that a 3-month intervention with soymilk reduced blood pressure in men and women with mild-to-moderate hypertension, and that this hypotensive effect was related to urinary excretion of the isoflavonoid genistein.⁴¹ Indeed, it has been shown that the BP-lowering effect of soy might be related to isoflavones,⁴² the suspected active ingredients in soy, via the activation of endothelial nitric oxide (NO) synthase (eNOS) and stimulation of NO production. Genistein is one of the soy isoflavones that can result in activation of eNOS and NO synthesis.⁴³ This is confirmed in another study where it was demonstrated that higher soy




Fig. 5. The funnel plot and the result of sensitivity analysis for SBP (A, B) and DBP (C, D).

consumption is related to higher plasma concentrations of NO.⁴⁴

It has been shown that oxidative stress and inflammation play a potential role in the development of hypertension⁴⁵; thereby suggesting that soy isoflavones might lower BP through antioxidant and anti-inflammatory effects.^{46,47} In line with the meta-analysis by Liu et al.,⁴⁸ it is unsurprising that we did not observe an inverse relationship of soy consumption with BP among older people, largely due to the adverse structural changes in the vessel wall among this group.⁵ It is known that BP increases with enhancing arterial stiffness, concomitant to aging, and the consumption of phytoestrogens have been inversely related to arterial stiffness.⁵ Moreover, soy consumption also reduces BP through its natriuretic effect, which is similar to furosemide.¹³

In the present study, we found that soy intake led to a higher reduction in blood pressure in participants with lower baseline systolic and diastolic BP, which is in contrast with result of the meta-analysis conducted by Liu et al.⁴⁸ One reason that should be noted for this inconstancy is the difference in inclusion criteria regarding study design, inter-study heterogeneity, and risk of bias, which was lower across our pooled included studies.

4.1. Strengths and limitations

The present meta-analysis is the first comprehensive study to have assessed the influence of soy intake on BP from randomized double-blind placebo-controlled trials. We used a comprehensive and accurate systematic search strategy, that allowed us to examine both indexed and non-indexed trials. To reduce between-study heterogeneity and

potential bias, and also enhance the power of results, we only included studies in which the control group received a placebo and where a double-blind design was employed. This approach improved quality of pooled analyses and subsequently permitting more rigorous insights in the effect of soy intake on BP. In addition, based on risk of bias assessment using Cochrane methodology, the quality of all studies was assessed overall as low risk. Subgroup analysis according to covariates that have clinical importance regarding soy intake and blood pressure improvements were pre-specified and provided additional insights to inform subsequent recommendations. It should be noted that there was no significant publication bias found, whilst pooled results were not sensitive to any individual study. Also, low amount of inter-study heterogeneity empowered the results. However, despite the clear novelty of this work, our study has some limitations that should be noted. First, as some included trials only reported dose of isoflavones, whilst others only reported soy, we were not able to apply a dose-response analysis. Second, we could not conduct subgroup analysis for baseline BMI and body weight of participants, which could be clinically important markers related to the change of BP, because of their absence in some included trials. Moreover, due to a variety of diseases among included study participants and lack of essential clinical data about their metabolic markers, we were not able to conduct additional subgroup analysis based on type of diseases and different health conditions without reducing the power of the analysis. Finally, although we contacted some corresponding authors to request missing essential data, in most cases, we received no response, or unsatisfactory responses.

4.2. Conclusion

In the present study, pooled effect sizes from 17 studies revealed a significant improvement in SBP and DBP in adults following soy consumption, in comparison with controls. In addition, subgroup analysis indicated a further reduction in both SBP and DBP in younger participants with lower baseline DBP and intervention durations < 16 weeks. Thus, increases to soy consumption could be considered as an alternative or complementary approach to improving BP outcomes among adults, and particularly among younger adults.

Authors contribution

All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published.

Funding

None.

Ethics approval and consent to participate

Not applicable.

Declaration of Competing Interest

The authors report no declarations of interest.

Acknowledgements

We thank Morteza Zare (https://www.researchgate.net/profile/Morteza_Zare), the expert statistician and head of Meta-Research Innovation office (METRIO), Shiraz University of Medical Sciences, Iran.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ctim.2021.102692>.

References

- Zeng J, Feng Y, Feng J, et al. The effect of soy intervention on insulin-like growth factor 1 levels: a meta-analysis of clinical trials. *Phytother Res*. 2020;34(July(7)): 1570–1577. <https://doi.org/10.1002/ptr.6630>. PubMed PMID: 32072706.
- Balla F. Nutritional value and economic aspects of fortification of foods of plant origin with soy protein. *J Am Oil Chem Soc*. 1974;51(January(1)):156A–158A. <https://doi.org/10.1007/BF02542121>. PubMed PMID: 4856014.
- Endres JG. *Soy protein products: Characteristics, nutritional aspects, and utilization*. The American Oil Chemists Society; 2001.
- Sugano M. *Soy in health and disease prevention*. CRC Press; 2005.
- Man B, Cui C, Zhang X, et al. The effect of soy isoflavones on arterial stiffness: a systematic review and meta-analysis of randomized controlled trials. *Eur J Nutr*. 2020;11(June):1–12. <https://doi.org/10.1007/s00394-020-02300-6>. PubMed PMID: 32529287.
- Schwingshackl L, Chaimani A, Schwedhelm C, et al. Comparative effects of different dietary approaches on blood pressure in hypertensive and pre-hypertensive patients: a systematic review and network meta-analysis. *Crit Rev Food Sci Nutr*. 2019;59(16): 2674–2687. <https://doi.org/10.1080/10408398.2018.1463967>. PubMed PMID: 29718689.
- Miller 3rd ER, Erlinger TP, Young DR, et al. Results of the Diet, Exercise, and Weight Loss I intervention Trial (DEW-IT). *Hypertension*. 2002;40(5):612–618.
- Schwingshackl L, Chaimani A, Hoffmann G, et al. Impact of different dietary approaches on blood pressure in hypertensive and prehypertensive patients: protocol for a systematic review and network meta-analysis. *BMJ Open*. 2017;7(April (4)): e014736. <https://doi.org/10.1136/bmjopen-2016-014736>. PubMed PMID: 28446526; PubMed Central PMCID: PMC5566592.
- Moore TJ, Conlin PR, Ard J, et al. DASH (Dietary Approaches to stop Hypertension) diet is effective treatment for stage 1 isolated systolic hypertension. *Hypertension*. 2001;38(2):155–158.
- Zhao L, Stamler J, Yan LL, et al. Blood pressure differences between northern and southern Chinese: role of dietary factors: the International Study on Macronutrients and Blood Pressure. *Hypertension*. 2004;43(6):1332–1337.
- Miller 3rd ER, Erlinger TP, Appel LJ. The effects of macronutrients on blood pressure and lipids: an overview of the DASH and OmniHeart trials. *Curr Atheroscler Rep*. 2006;8(November(6)):460–465. <https://doi.org/10.1007/s11883-006-0020-1>. PubMed PMID: 17045071.
- Katherine L, Ziegler TR. *Modern nutrition in health and disease*. 2012.
- Yang G, Shu X-O, Jin F, et al. Longitudinal study of soy food intake and blood pressure among middle-aged and elderly Chinese women. *Am J Clin Nutr*. 2005;81(5):1012–1017.
- Nagata C, Shimizu H, Takami R, et al. Association of blood pressure with intake of soy products and other food groups in Japanese men and women. *Prev Med*. 2003;36(June(6)):692–697. [https://doi.org/10.1016/s0091-7435\(03\)00052-5](https://doi.org/10.1016/s0091-7435(03)00052-5). PubMed PMID: 12744912.
- Maleki Z, Jazayeri S, Eslami O, et al. Effect of soy milk consumption on glycemic status, blood pressure, fibrinogen and malondialdehyde in patients with non-alcoholic fatty liver disease: a randomized controlled trial. *Complement Ther Med*. 2019;44(June):44–50. <https://doi.org/10.1016/j.ctim.2019.02.020>. PubMed PMID: 31126574; eng.
- Sedaghat A, Shahbazian H, Rezazadeh A, et al. The effect of soy nut on serum total antioxidant, endothelial function and cardiovascular risk factors in patients with type 2 diabetes. *Diabetes Metab Syndr*. 2019;13(Mar-Apr(2)):1387–1391. <https://doi.org/10.1016/j.dsx.2019.01.057>. PubMed PMID: 31336497; eng.
- Kim J, Lee H, Lee O, et al. Isoflavone supplementation influenced levels of triglyceride and luteinizing hormone in Korean postmenopausal women. *Arch Pharm Res*. 2013;36(March(3)):306–313. <https://doi.org/10.1007/s12272-013-0059-9>. PubMed PMID: 23475289; eng.
- Wong WW, Taylor AA, Smith EO, et al. Effect of soy isoflavone supplementation on nitric oxide metabolism and blood pressure in menopausal women. *Am J Clin Nutr*. 2012;95(June(6)):1487–1494. <https://doi.org/10.3945/ajcn.111.032045>. PubMed PMID: 22552034; PubMed Central PMCID: PMC3349458. eng.
- Liu ZM, Ho SC, Chen YM, et al. Effect of whole soy and purified daidzein on ambulatory blood pressure and endothelial function—a 6-month double-blind, randomized controlled trial among Chinese postmenopausal women with prehypertension. *Eur J Clin Nutr*. 2015;69(October(10)):1161–1168. <https://doi.org/10.1038/ejcn.2015.24>. PubMed PMID: 25782428; eng.
- Husain D, Khanna K, Puri S, et al. Supplementation of soy isoflavones improved sex hormones, blood pressure, and postmenopausal symptoms. *J Am Coll Nutr*. 2015;34(1):42–48. <https://doi.org/10.1080/07315724.2013.875434>. PubMed PMID: 25648211; eng.
- He J, Gu D, Wu X, et al. Effect of soybean protein on blood pressure: a randomized, controlled trial. *Ann Intern Med*. 2005;143(July (1)):1–9. <https://doi.org/10.7326/0003-4819-143-1-200507050-00004>. PubMed PMID: 15998749; eng.
- Chan YH, Lau KK, Yiu KH, et al. Reduction of C-reactive protein with isoflavone supplement reverses endothelial dysfunction in patients with ischaemic stroke. *Eur Heart J*. 2008;29(November(22)):2800–2807. <https://doi.org/10.1093/eurheartj/ehn409>. PubMed PMID: 18812325.
- Sagara M, Kanda T, M NJ, et al. Effects of dietary intake of soy protein and isoflavones on cardiovascular disease risk factors in high risk, middle-aged men in Scotland. *J Am Coll Nutr*. 2004;23(February(1)):85–91. <https://doi.org/10.1080/07315724.2004.10719347>. PubMed PMID: 14963058; eng.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;15(4):264–269.
- Follmann D, Elliott P, Suh I, et al. Variance imputation for overviews of clinical trials with continuous response. *J Clin Epidemiol*. 1992;45(July(7)):769–773. [https://doi.org/10.1016/0895-4356\(92\)90054-q](https://doi.org/10.1016/0895-4356(92)90054-q). PubMed PMID: 1619456.
- Higgins JP, Altman DG, Gotzsche PC, et al. *The cochrane collaboration's tool for assessing risk of bias in randomised trials*. 343. 2011. d5928.
- Higgins J, Thomas J, Chandler J, et al. *Cochrane handbook for systematic reviews of interventions version 6.0 (updated July 2019)*. Cochrane. 2019.
- Borenstein M, Hedges LV, Higgins JP, et al. *Introduction to meta-analysis*. John Wiley & Sons; 2011.
- Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics*. 1994;50(December(4)):1088–1101. PubMed PMID: 7786990.
- Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(September (7109)):629–634. <https://doi.org/10.1136/bmj.315.7109.629>. PubMed PMID: 9310563; PubMed Central PMCID: PMC12127453.
- Teede HJ, Dalais FS, Kotsopoulos D, et al. Dietary soy has both beneficial and potentially adverse cardiovascular effects: a placebo-controlled study in men and postmenopausal women. *J Clin Endocrinol Metab*. 2001;86(July(7)):3053–3060. <https://doi.org/10.1210/jcem.86.7.7645>. PubMed PMID: 11443167; eng.
- Kim JI, Kim JC, Kang MJ, et al. Effects of pinitol isolated from soybeans on glycaemic control and cardiovascular risk factors in Korean patients with type II diabetes mellitus: a randomized controlled study. *Eur J Clin Nutr*. 2005;59(March(3)): 456–458. <https://doi.org/10.1038/sj.ejcn.1602081>. PubMed PMID: 15536472; eng.
- Kwak JH, Lee JH, Ahn CW, et al. Black soy peptide supplementation improves glucose control in subjects with prediabetes and newly diagnosed type 2 diabetes mellitus. *J Med Food*. 2010;13(December(6)):1307–1312. <https://doi.org/10.1089/jmf.2010.1075>. PubMed PMID: 21091245.
- Liu ZM, Ho SC, Chen YM, et al. Effect of soy protein and isoflavones on blood pressure and endothelial cytokines: a 6-month randomized controlled trial among postmenopausal women. *J Hypertens*. 2013;31(February(2)):384–392. <https://doi.org/10.1097/HJH.0b013e32835c0905>. PubMed PMID: 23203140; eng.

- 35 Cheng WC, Lo SC, Tsai KS, et al. Effects of high-dose phytoestrogens on circulating cellular microparticles and coagulation function in postmenopausal women. *J Formos Med Assoc.* 2015;114(August(8)):710–716. <https://doi.org/10.1016/j.jfma.2013.11.001>. PubMed PMID: 24360978; eng.
- 36 Hermansen K, Hansen B, Jacobsen R, et al. Effects of soy supplementation on blood lipids and arterial function in hypercholesterolaemic subjects. *Eur J Clin Nutr.* 2005;59(July(7)):843–850. <https://doi.org/10.1038/sj.ejcn.1602151>. PubMed PMID: 15900307; eng.
- 37 Squadrito F, Marini H, Bitto A, et al. Genistein in the metabolic syndrome: results of a randomized clinical trial. *J Clin Endocrinol Metab.* 2013;98(Augu (8)):3366–3374. <https://doi.org/10.1210/jc.2013-1180>. PubMed PMID: 23824420.
- 38 Aubertin-Leheudre M, Lord C, Khalil A, et al. Isoflavones and clinical cardiovascular risk factors in obese postmenopausal women: a randomized double-blind placebo-controlled trial. *J Womens Health (Larchmt).* 2008;17(October(8)):1363–1369. <https://doi.org/10.1089/jwh.2008.0836>. PubMed PMID: 18788990.
- 39 Washburn S, Burke GL, Morgan T, et al. Effect of soy protein supplementation on serum lipoproteins, blood pressure, and menopausal symptoms in perimenopausal women. *Menopause.* 1999;6(1):7–13. Spring PubMed PMID: 10100174.
- 40 Welty FK, Lee KS, Lew NS, et al. Effect of soy nuts on blood pressure and lipid levels in hypertensive, prehypertensive, and normotensive postmenopausal women. *Arch Intern Med.* 2007;167(May(10)):1060–1067. <https://doi.org/10.1001/archinte.167.10.1060>. PubMed PMID: 17533209.
- 41 Rivas M, Garay RP, Escanero JF, et al. Soy milk lowers blood pressure in men and women with mild to moderate essential hypertension. *J Nutr.* 2002;132(July(7)):1900–1902. <https://doi.org/10.1093/jn/132.7.1900>. PubMed PMID: 12097666; eng.
- 42 Zarei A, Stasi C, Mahmoodi M, et al. Effect of soy consumption on liver enzymes, lipid profile, anthropometry indices, and oxidative stress in patients with non-alcoholic fatty liver disease: a systematic review and meta-analysis of clinical trials. *J Iranian J Basic Med Sci.* 2020;23(10):1245–1250. <https://doi.org/10.22038/ijbms.2020.46854.10797>.
- 43 Richardson SI, Steffen LM, Swett K, et al. Dietary total isoflavone intake is associated with lower systolic blood pressure: The Coronary Artery Risk Development in Young Adults (CARDIA) study. *J Clin Hypertens.* 2016;18(8):778–783.
- 44 Hallund J, Bugel S, Tholstrup T, et al. Soya isoflavone-enriched cereal bars affect markers of endothelial function in postmenopausal women. *Br J Nutr.* 2006;95(June (6)):1120–1126. <https://doi.org/10.1079/bjn20061734>. PubMed PMID: 16768834.
- 45 Jalali M, Karamizadeh M, Ferns GA, et al. The effects of cashew nut intake on lipid profile and blood pressure: a systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med.* 2020;50(May):102387. <https://doi.org/10.1016/j.ctim.2020.102387>. PubMed PMID: 32444052.
- 46 Panneerselvam S, Packirisamy R, Bobby Z, et al. Protective effect of soy isoflavones (from Glycine max) on adipose tissue oxidative stress and inflammatory response in an experimental model of post-menopausal obesity: the molecular mechanisms. *Biochem Anal Biochem.* 2016;5(266), 2161-1009.1000266.
- 47 Yoon GA, Park S. Antioxidant action of soy isoflavones on oxidative stress and antioxidant enzyme activities in exercised rats. *Nutr Res Pract.* 2014;8(December(6)): 618–624. <https://doi.org/10.4162/nrp.2014.8.6.618>. PubMed PMID: 25489400; PubMed Central PMCID: PMC4252520.
- 48 Liu XX, Li SH, Chen JZ, et al. Effect of soy isoflavones on blood pressure: a meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis.* 2012;22(June (6)):463–470. <https://doi.org/10.1016/j.numecd.2010.09.006>. PubMed PMID: 21310599.
- 49 Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to meta-analysis.* John Wiley & Sons; 2011.
- 50 Follmann D, Elliott P, Suh IL, Cutler J. Variance imputation for overviews of clinical trials with continuous response. *J clin epidemiol.* 1992;45(7):769–773.